

## Dewormer Chart for Camelids

Please read notes below before using this chart

CAMELID CHART 2015	Valbazen® (albendazole)	Safe-Guard® or Panacur® (fenbendazole)	Ivomec® Sheep Drench (ivermectin)	Cydectin® Sheep Drench (moxidectin)	Levasole®, Prohibit® Tramisol® (levamisole)
Weight Pounds	DOSE: 20 mg/kg orally	DOSE: 20 mg/kg orally	DOSE: 0.4 mg/kg orally	DOSE: 0.4 mg/kg orally	DOSE: 8 mg/kg orally
	Dose in mls	Dose in mls	Dose in mls	Dose in mls	Dose in mls*
10	0.8	0.9	2.3	1.8	0.8
20	1.6	1.8	4.5	3.6	1.6
30	2.4	2.7	6.8	5.4	2.4
40	3.2	3.6	9.0	7.3	3.3
<b>50</b>	<b>4.0</b>	<b>4.5</b>	<b>11.4</b>	<b>9.0</b>	<b>4.1</b>
60	4.8	5.5	13.6	11	4.9
70	5.6	6.4	16	12.7	5.7
80	6.4	7.3	18.2	14.6	6.5
90	7.2	8.2	21	16.4	7.3
<b>100</b>	<b>8.0</b>	<b>9.1</b>	<b>22.7</b>	<b>18</b>	<b>8.1</b>
110	8.8	10	25	20	8.9
120	9.6	10.9	27.3	22	9.7
130	10.4	11.8	29.5	23.6	10.6
140	11.2	12.7	31.8	25.4	11.4
<b>150</b>	<b>12.0</b>	<b>13.6</b>	<b>34</b>	<b>27.3</b>	<b>12.2</b>
160	12.8	14.5	36.4	29.1	13.0
170	13.6	15.5	38.6	31	13.8
180	14.4	16.4	42	33	14.6
190	15.2	17.3	43.2	34.6	15.5
<b>200</b>	<b>16</b>	<b>18.2</b>	<b>45.5</b>	<b>36.4</b>	<b>16.2</b>
210	16.8	19.1	47.7	38.2	17.1
220	17.6	20	50	40	17.9
230	18.4	20.9	52.3	41.8	18.7
240	19.2	21.8	54.5	43.6	19.5
<b>250</b>	<b>20</b>	<b>22.7</b>	<b>56.8</b>	<b>45.4</b>	<b>20.3</b>
260	20.8	23.6	59	47.3	21.2
270	21.6	24.5	61.3	49.1	22.0
280	22.4	25.5	63.6	50.9	22.8
290	23.2	26.4	66	52.7	23.6
<b>300</b>	<b>24</b>	<b>27.3</b>	<b>68.2</b>	<b>54.6</b>	<b>24.4</b>
310	24.8	28.2	70.4	56.4	25.2
320	25.6	29.1	72.7	58.2	26.0
330	26.4	30	75	60	26.8
340	27.2	30.9	77.3	61.8	27.7
<b>350</b>	<b>28</b>	<b>31.8</b>	<b>79.5</b>	<b>63.6</b>	<b>28.5</b>
360	28.8	32.7	81.8	65.4	29.3

\***Levamisole:** use dosing recommendations indicated in this chart *only* if you dilute the levamisole soluble drench powder in this manner: dissolve contents of 13 gm (11.7 gm active ingredient) packet in 262 ml water. A 52-gram packet is dissolved in 1048 ml water (44.7 mg/ml).

**Albendazole:** do not use in first 30 days of pregnancy. **Do not** re-dose on consecutive days.

Lisa Williamson, DVM, MS; University of Georgia College of Veterinary Medicine, Athens, Georgia 30602